

Regional School Unit 57

86 West Road, Waterboro, Maine 04087

Tel. No. (207) 247-3221

Fax. No. (207) 247-3477

Larry Malone
Superintendent

Colin M. Walsh, CPA
Director of Finance and Operations

Lori Lodge
Curriculum Coordinator

Susan Prince
Director of Special Education

Nicole Poole
Asst. Director of Special Education

RSU 57 prepares respectful, responsible and creative thinkers for success in the global community.

June 01, 2017


Dear MHS Parents/Guardians,

The Maine Center for Disease Control and Prevention (Maine CDC) has notified Massabesic High School of a confirmed case of Pertussis, more commonly known as whooping cough. Pertussis is most common in adolescents and has been reported in all of the counties in Maine. The MHS Nurse is working directly with the Maine CDC regarding this case and will provide further notification if appropriate.

There is a chance your child may have been exposed to Pertussis. The Maine CDC is recommending that only students exhibiting symptoms, particularly a severe cough that causes vomiting and/or gagging, or a cough lasting 2 weeks or more, be tested for the illness. Contact your child's physician with concerns. Once confirmed, Pertussis is treated with a 5-day course of antibiotics. Massabesic High School has been reassured that the affected student has been prescribed appropriate treatment and should no longer be infectious to others.

The Maine CDC recommends all children aged 11-18 receive the Tdap booster. Please see the attached fact sheet for more information. If you have any further questions or concerns, please contact the MHS Nurse, Nancy Sobanik at 247-3141, ext. 2215, or the Maine CDC at 1-800-821-5821.

Thank you,



Larry Malone
Superintendent

Pertussis

Fact Sheet



Maine Center for Disease
Control and Prevention

An Office of the
Department of Health and Human Services

Paul R. LePage, Governor

Mary C. Mayhew, Commissioner

What is pertussis?

Pertussis, or whooping cough, is caused by bacteria called *Bordetella pertussis*. Pertussis can be very serious, especially in infants. It mainly affects the breathing.

What are the signs of pertussis?

The first signs of pertussis are similar to a cold (sneezing, runny nose, low-grade fever, and a cough). These signs start 5 to 21 days after exposure. After one or two weeks of illness, the cough gets worse with symptoms that may include:

- A sudden, uncontrollable cough where one cough follows the next without a break for breath.
- A high-pitched whooping sound when breathing in after a coughing episode. Whooping is less common in infants and adults.
- Vomiting after coughing
- Exhaustion

Over time, coughing spells become less frequent, but may continue for several weeks or months.

How do you catch pertussis?

Pertussis is spread from person to person through the air. A person has to breathe in droplets from an infected person to get sick. For example, a person may catch pertussis by standing close (less than 3 feet away) to an infected person who is coughing or sneezing.

When and for how long can a person spread pertussis?

Pertussis is most likely to spread to others early in the illness. Persons with pertussis can no longer spread the disease once they have completed 5 days of treatment with antibiotics. However, persons with pertussis who do not take antibiotics can spread the disease during the first 21 days they are sick.

What are the complications?

Severe pertussis is more likely in infants than in older children or adults. The most common complication of pertussis is pneumonia. Seizures, swelling of the brain and death are rare but possible.

Who gets pertussis?

People of all ages can get pertussis, even people who have been vaccinated or have had pertussis infection in the past. It is most common in school-aged children and teenagers, but cases in adults also occur. Older children and adults usually have less severe illness, but they can still spread the disease to infants and young children.

Is there any treatment?

Pertussis can be treated with antibiotics. Treatment should be started early to prevent the spread to others and can lessen the symptoms of illness. A cough may continue for many weeks after treatment especially if treatment wasn't started until late in the illness. Pertussis bacteria die off naturally after three weeks of coughing. Antibiotics are not recommended if a person has already been coughing for 21 days or more.

Why did my healthcare provider tell me or my child to stay home for 5 days?

You can still spread pertussis to others until you take 5 days of antibiotics. Disease can spread quickly in schools or the workplace.

What will happen if I do not want to take the antibiotics?

If you do not take the antibiotics, you will need to stay away from others for 21 days after you start to become sick. This includes staying home from daycare, school, work or events.

Is there a lab test for pertussis?

Yes. To test for pertussis, your healthcare provider

may insert a swab (like a long q-tip) into your nose. The lab will test the material on the swab to see if they can find the bacteria that causes pertussis.

[ase/epi/index.shtml](#)

The federal Centers for Disease Control and Prevention website - <http://www.cdc.gov> – is another excellent source of health information.

Is there a vaccine for pertussis?

Yes, there are vaccines. The childhood vaccine is called Dtap, and the Pertussis booster for adolescents and adults is called Tdap.

People who catch pertussis after being vaccinated have milder illness and are much less likely to be hospitalized or die from the disease.

Vaccines prevent 200,000 cases of pertussis each year in the US. Talk to your healthcare provider about getting vaccinated against pertussis.

How can pertussis be prevented?

Vaccinate all children on time. Speak with your healthcare provider to see if you should get a booster vaccine. This is the best way to prevent pertussis.

Other ways to prevent pertussis are to:

- Avoid close contact with others who are sick or coughing
- Wash your hands often
- Stay at home if you are ill
- Cover your cough with a tissue or cough into your sleeve
- Use routine cleaners and disinfectants to remove these bacteria from surfaces or objects.
- See your healthcare provider if you have signs of pertussis or have been in close contact with someone who has pertussis.

Where can I get more information?

For more information contact your healthcare provider or local health center. Infectious Disease Epidemiology website:

<http://www.maine.gov/dhhs/mecdc/infectious-dise>